

ZSCAN2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12089**Specification**

ZSCAN2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q7Z7L9
Other Accession	NM_181877 , NP_870992
Reactivity	Human, Mouse, Rat, Pig, Bovine, Dog
Predicted	Human, Mouse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67kDa KDa

ZSCAN2 antibody - N-terminal region - Additional Information**Gene ID** 54993**Alias Symbol** ZFP29, ZNF854**Other Names**

Zinc finger and SCAN domain-containing protein 2, Zinc finger protein 29 homolog, Zfp-29, Zinc finger protein 854, ZSCAN2, ZFP29, ZNF854

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZSCAN2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZSCAN2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZSCAN2 antibody - N-terminal region - Protein Information**Name** ZSCAN2**Synonyms** ZFP29, ZNF854**Function**

May be involved in transcriptional regulation during the post-meiotic stages of spermatogenesis.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.

ZSCAN2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZSCAN2 antibody - N-terminal region - Images



WB Suggested Anti-ZSCAN2 Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate

ZSCAN2 antibody - N-terminal region - References

Denny, P. (1991) Gene 106(2), 221-227. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.