

ANK1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12117**Specification**

ANK1 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q53ER1
Other Accession	NM_020478 , NP_065211
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Horse, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17kDa KDa

ANK1 antibody - C-terminal region - Additional Information

Alias Symbol **ANK, SPH1, SPH2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ANK1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ANK1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ANK1 antibody - C-terminal region - Protein Information**ANK1 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ANK1 antibody - C-terminal region - Images



Human Skin



Human Lung



WB Suggested Anti-ANK1 Antibody Titration: 1.25µg/ml

Positive Control: Jurkat cell lysate

ANK1 antibody - C-terminal region - References

Chang, T.L., (2004) Cell. Mol. Biol. (Noisy-le-grand) 50(1), 59-66 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycle S.