

GH2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12127**Specification**

GH2 antibody - middle region - Product Information

Application	WB
Primary Accession	O14644
Other Accession	NM_022558 , NP_072052
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27kDa KDa

GH2 antibody - middle region - Additional Information

Alias Symbol **GH-V, GHL, GHV, hGH-V**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GH2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


GH2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

GH2 antibody - middle region - Protein Information**GH2 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GH2 antibody - middle region - Images



90 kDa
65 kDa
40 kDa
29 kDa
22 kDa

WB Suggested Anti-GH2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: PANC1 cell lysate

GH2 is supported by BioGPS gene expression data to be expressed in PANC1

GH2 antibody - middle region - References

0ReconstitutionandStorage:Forshorttermuse,storeat2-8Cupto1week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.