

GAMT antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12141**Specification**

GAMT antibody - middle region - Product Information

Application	IHC, WB
Primary Accession	A8K0A0
Other Accession	NM_138924 , NP_620279
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29kDa KDa

GAMT antibody - middle region - Additional Information

Alias Symbol **PIG2, TP53I2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GAMT antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


GAMT antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

GAMT antibody - middle region - Protein Information**GAMT antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GAMT antibody - middle region - Images



Rabbit Anti-GAMT Antibody

Formalin Fixed Paraffin Embedded Tissue: Human Adult liver Observed Staining: Cytoplasmic

Primary Antibody

Concentration: 1:100

Secondary Antibody: Donkey anti-Rabbit-Cy2/3


Secondary Antibody

Concentration: 1:200

Magnification: 20X

Exposure Time: 0.5 – 2.0 sec

Protocol located in Reviews and Data.



WB Suggested Anti-GAMT Antibody Titration: 0.2-1 µg/ml

Positive Control: Human Thymus

GAMT antibody - middle region - References

Almeida, L.S., (2007) Mol. Genet. Metab. 91(1), 1-6 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.