

Parp8 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12176**Specification**

Parp8 antibody - C-terminal region - Product Information

Application	WB
Other Accession	XM_978094 , XP_983188
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

Parp8 antibody - C-terminal region - Additional Information

Alias Symbol **281043008Rik, D13Ertd275e**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Parp8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


Parp8 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Parp8 antibody - C-terminal region - Protein Information**Parp8 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Parp8 antibody - C-terminal region - Images



WB Suggested Anti-Parp8 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Mouse Thymus