

TJP2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12188**Specification**

TJP2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A2A3H9
Other Accession	NM_201629 , NP_963923
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	118kDa KDa

TJP2 antibody - C-terminal region - Additional Information

Alias Symbol	MGC26306, X104, ZO-2, ZO2, DFNA51, DUP9q21.11, C9DUPq21.11
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TJP2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


TJP2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TJP2 antibody - C-terminal region - Protein Information**TJP2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TJP2 antibody - C-terminal region - Images



WB Suggested Anti-TJP2 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

TJP2 antibody - C-terminal region - References

Bel(2008)Hum.Genet.123(3),267-272ReconstitutionandStorage:Forshorttermuse,storeat2-8Cupto1 week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.