

VPS41 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12227**Specification**

VPS41 antibody - middle region - Product Information

Application	WB
Primary Accession	P49754
Other Accession	NM_080631 , NP_542198
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	96kDa KDa

VPS41 antibody - middle region - Additional Information**Gene ID** 27072

Alias Symbol	HVPS41, HVSP41, hVps41p
Other Names	
Vacuolar protein sorting-associated protein 41 homolog, S53, VPS41	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-VPS41 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

VPS41 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

VPS41 antibody - middle region - Protein Information**Name** VPS41**Function**

Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Acts as a component of the HOPS endosomal tethering complex. This complex is proposed to be involved in the Rab5- to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic towards lysosomes (PubMed:23351085, PubMed:33851776). Involved in homotypic vesicle fusions between late endosomes and in heterotypic fusions between late endosomes and lysosomes implicated in degradation of endocytosed cargo (PubMed:23167963, PubMed:25445562, PubMed:25908847, PubMed:9159129). Required for fusion of autophagosomes with lysosomes (PubMed:25783203, PubMed:37821429). Links the HOPS complex to endosomal Rab7 via its association with RILP and to lysosomal membranes via its association with ARL8B, suggesting that these interactions may bring the compartments to close proximity for fusion (PubMed:21802320, PubMed:25445562, PubMed:25908847). Involved in the direct trans-Golgi network to late endosomes transport of lysosomal membrane proteins independently of HOPS (PubMed:23322049). Involved in sorting to the regulated secretory pathway presumably implicating the AP-3 adapter complex (By similarity). May play a role in HOPS-independent function in the regulated secretory pathway (PubMed:24210660).

Cellular Location

Endosome membrane; Peripheral membrane protein. Late endosome membrane; Peripheral membrane protein. Early endosome membrane; Peripheral membrane protein. Lysosome membrane; Peripheral membrane protein. Golgi apparatus, trans- Golgi network. Cytoplasmic vesicle, clathrin-coated vesicle. Cytoplasm, cytosol

Tissue Location


Expressed in cerebral cortex and cerebellum. Highly expressed in Purkinje cells.


VPS41 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VPS41 antibody - middle region - Images





WB Suggested Anti-VPS41 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate