

HERC6 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12243**Specification**

HERC6 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8IVU3
Other Accession	NM_017912 , NP_060382
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Rabbit, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	115kDa KDa

HERC6 antibody - N-terminal region - Additional Information**Gene ID** 55008**Alias Symbol** **FLJ20637****Other Names**

Probable E3 ubiquitin-protein ligase HERC6, 6.3.2.-, HECT domain and RCC1-like domain-containing protein 6, HERC6

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-HERC6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

HERC6 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HERC6 antibody - N-terminal region - Protein Information**Name** HERC6**Function**

E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates.

Cellular Location

Cytoplasm, cytosol.


Tissue Location

Detected in brain, heart, placenta and testis.

HERC6 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HERC6 antibody - N-terminal region - Images

WB Suggested Anti-HERC6 Antibody Titration: 0.2-1 μ g/ml
Positive Control: ACHN cell lysate