

DTX2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12253**Specification**

DTX2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6XM87
Other Accession	NM_001102596 , NP_001096066
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Goat, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Pig, Chicken, Goat, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62kDa kDa

DTX2 antibody - C-terminal region - Additional Information

Alias Symbol **KIAA1528, MGC71098, RNF58**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DTX2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


DTX2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DTX2 antibody - C-terminal region - Protein Information**DTX2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DTX2 antibody - C-terminal region - Images



WB Suggested Anti-DTX2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: HepG2 cell lysate