

RNF39 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12278**Specification**

RNF39 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A6NCD6
Other Accession	NM_170770 , NP_739576
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28kDa KDa

RNF39 antibody - C-terminal region - Additional Information

Alias Symbol **HZF, HZFW, LIRF**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-RNF39 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


RNF39 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RNF39 antibody - C-terminal region - Protein Information**RNF39 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RNF39 antibody - C-terminal region - Images



WB Suggested Anti-RNF39 Antibody Titration: 2.5µg/ml
Positive Control: HepG2 cell lysate

RNF39 antibody - C-terminal region - References

Matsuo, R., (2001) Biochem. Biophys. Res. Commun. 289(2), 479-484 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.