

Abcf1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12306**Specification**

Abcf1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6MG08
Other Accession	NM_001109883 , NP_001103353
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	95kDa KDa

Abcf1 antibody - C-terminal region - Additional Information**Gene ID** 85493**Alias Symbol** **Abc50, Abcf1****Other Names**

ATP-binding cassette sub-family F member 1, ATP-binding cassette 50, Abcf1, Abc50

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Abcf1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Abcf1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Abcf1 antibody - C-terminal region - Protein Information**Name** Abcf1**Synonyms** Abc50**Function**

Required for efficient Cap- and IRES-mediated mRNA translation initiation. Not involved in the ribosome biogenesis (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q8NE71}. Nucleus, nucleoplasm


{ECO:0000250|UniProtKB:Q8NE71}. Nucleus envelope {ECO:0000250|UniProtKB:Q8NE71}

Abcf1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Abcf1 antibody - C-terminal region - Images



Researcher: Dr. Joanna Stewart, Centre for Biological Sciences, University of Southampton
Application: IP

Species+tissue/cell type: 1: 500 ug HEK293 cell lysate transfected with ABCF1 wt, 2: 500 ug HEK293 cell lysate untransfected, 3: 500 ug HEK293 cell lysate transfected with mutABCF1
Amount of IP antibody: 1ug

Primary Antibody Dilution: 1:1000

Secondary Antibody: Anti-rabbit-Alexa Fluor 680

Secondary Antibody Dilution: 1:20000