

SLC10A5 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12318**Specification**

SLC10A5 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q5PT55
Other Accession	NM_001010893 , NP_001010893
Reactivity	Human, Rat, Pig, Horse
Predicted	Rat, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

SLC10A5 antibody - C-terminal region - Additional Information**Gene ID** 347051**Alias Symbol** P5**Other Names**

Sodium/bile acid cotransporter 5, Na(+)/bile acid cotransporter 5, Solute carrier family 10 member 5, SLC10A5

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-SLC10A5 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SLC10A5 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SLC10A5 antibody - C-terminal region - Protein Information**Name** SLC10A5**Cellular Location**


Membrane; Multi-pass membrane protein

SLC10A5 antibody - C-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC10A5 antibody - C-terminal region - Images



Human Lung



WB Suggested Anti-SLC10A5 Antibody Titration: 1.25µg/ml
Positive Control: Jurkat cell lysate

SLC10A5 antibody - C-terminal region - References

Ota, T., (2004) Nat. Genet. 36(1), 40-45 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.