

**SLC35C1 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI12361****Specification**

---

**SLC35C1 antibody - N-terminal region - Product Information**

|                   |                                                                    |
|-------------------|--------------------------------------------------------------------|
| Application       | WB                                                                 |
| Primary Accession | <a href="#">O96A29</a>                                             |
| Other Accession   | <a href="#">NM_018389</a> , <a href="#">NP_060859</a>              |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog     |
| Predicted         | Human, Mouse, Rat, Rabbit, Chicken, Horse, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                                             |
| Clonality         | Polyclonal                                                         |
| Calculated MW     | 40kDa KDa                                                          |

**SLC35C1 antibody - N-terminal region - Additional Information****Gene ID** 55343**Alias Symbol** **FLJ11320, FLJ14841, FUCT1, CDG2C**  
**Other Names**  
GDP-fucose transporter 1, Solute carrier family 35 member C1, SLC35C1, FUCT1**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 100 ul of distilled water. Final anti-SLC35C1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

SLC35C1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**SLC35C1 antibody - N-terminal region - Protein Information****Name** SLC35C1 ([HGNC:20197](#))**Function**

Antiporter specific for GDP-l-fucose and depending on the concomitant reverse transport of GMP. Involved in GDP-fucose import from the cytoplasm into the Golgi lumen.

**Cellular Location**


Golgi apparatus membrane; Multi-pass membrane protein

## SLC35C1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## SLC35C1 antibody - N-terminal region - Images



WB Suggested Anti-SLC35C1 Antibody Titration: 1.25µg/ml  
Positive Control: Jurkat cell lysate

## SLC35C1 antibody - N-terminal region - References

Hidalgo, A., (2003) Blood 101(5), 1705-1712 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.