

SLC12A8 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12366**Specification**

SLC12A8 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	A0AV02
Other Accession	NM_024628 , NP_078904
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78kDa KDa

SLC12A8 antibody - N-terminal region - Additional Information**Gene ID** 84561**Alias Symbol** CCC9, DKFZp686L18248, FLJ23188**Other Names**

Solute carrier family 12 member 8, Cation-chloride cotransporter 9, SLC12A8, CCC9

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SLC12A8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SLC12A8 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SLC12A8 antibody - N-terminal region - Protein Information**Name** SLC12A8**Synonyms** CCC9**Function**

Cation/chloride cotransporter that may play a role in the control of keratinocyte proliferation.

Cellular Location

Membrane; Multi-pass membrane protein


Tissue Location

Ubiquitous with very low level in normal skin.

SLC12A8 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC12A8 antibody - N-terminal region - Images

WB Suggested Anti-SLC12A8 Antibody Titration: 0.2-1 µg/ml
Positive Control: THP-1 cell lysate