

SLC2A10 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12369**Specification**

SLC2A10 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O95528
Other Accession	NM_030777 , NP_110404
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa KDa

SLC2A10 Antibody - C-terminal region - Additional Information**Gene ID** 81031**Alias Symbol** **ATS, GLUT10****Other Names**

Solute carrier family 2, facilitated glucose transporter member 10, Glucose transporter type 10, GLUT-10, SLC2A10, GLUT10

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SLC2A10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SLC2A10 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SLC2A10 Antibody - C-terminal region - Protein Information**Name** SLC2A10 {ECO:0000303|PubMed:11247674, ECO:0000312|HGNC:HGNC:13444}**Function**

Facilitative glucose transporter required for the development of the cardiovascular system.

Cellular Location

Endomembrane system; Multi-pass membrane protein. Cytoplasm, perinuclear region

Tissue Location


Widely expressed; highest levels in liver and pancreas.

SLC2A10 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC2A10 Antibody - C-terminal region - Images



Host: Rabbit

Target Name: SLC2A10

Sample Tissue: NCI-H226 Whole cell lysate

s

Antibody Dilution: 1.0µg/ml