

MICA antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12395**Specification**

MICA antibody - middle region - Product Information

Application	WB
Primary Accession	Q29983.1
Other Accession	NM_000247 , NP_000238
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

MICA antibody - middle region - Additional Information

Alias Symbol **DAQB-48K1.7, MGC111087, PERB11.1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-MICA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


MICA antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

MICA antibody - middle region - Protein Information**MICA antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MICA antibody - middle region - Images



70 kDa
60 kDa
48 kDa
36 kDa
21 kDa

WB Suggested Anti-MICA Antibody Titration: 2.5µg/ml
Positive Control: Jurkat cell lysate

MICA antibody - middle region - References

Boissel, N., (2006) J. Immunol. 176(8), 5108-5116 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.