

Ppic Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12412**Specification**

Ppic Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P30412
Other Accession	NM_008908 , NP_032934
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Mouse, Rat, Pig, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 23kDa KDa

Ppic Antibody - C-terminal region - Additional Information**Gene ID** 19038

Alias Symbol	CyP-20°C
Other Names	
Peptidyl-prolyl cis-trans isomerase C, PPlase C, 5.2.1.8, Cyclophilin C, Rotamase C, Ppic, Cypc	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ppic antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ppic Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ppic Antibody - C-terminal region - Protein Information**Name** Ppic {ECO:0000312|MGI:MGI:97751}**Function**

PPlase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding.

Cellular Location

Cytoplasm.

Tissue Location


Specifically expressed in kidney.

Ppic Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ppic Antibody - C-terminal region - Images



WB Suggested Anti-Ppic Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Heart