

OR6C68 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12435**Specification**

OR6C68 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	A6NDL8
Other Accession	NM_001005519 , NP_001005519
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36kDa KDa

OR6C68 antibody - N-terminal region - Additional Information**Gene ID** 403284**Other Names**

Olfactory receptor 6C68, OR6C68

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-OR6C68 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

OR6C68 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

OR6C68 antibody - N-terminal region - Protein Information**Name** OR6C68**Function**

Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR6C68 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR6C68 antibody - N-terminal region - Images



WB Suggested Anti-OR6C68 Antibody Titration: 0.2-1 μ g/ml
Positive Control: OVCAR-3 cell lysate