

WDR33 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12440**Specification**

WDR33 antibody - middle region - Product Information

Application	WB
Primary Accession	Q6NUQ0
Other Accession	NM_001006623 , NP_001006624
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Zebrafish, Pig, Chicken, Horse, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30kDa KDa

WDR33 antibody - middle region - Additional Information

Alias Symbol **FLJ11294, WDC146, NET14**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-WDR33 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


WDR33 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

WDR33 antibody - middle region - Protein Information**WDR33 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WDR33 antibody - middle region - Images




WB Suggested Anti-WDR33 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

WDR33 is supported by BioGPS gene expression data to be expressed in HepG2



Lanes: 1: SREC pulldown from lysate from 10⁶ human 293T cells


Primary Antibody Dilution: 1:1000

Secondary Antibody: Anti-rabbit-Alexa Fluor

Secondary Antibody Dilution: 1:5000

Gene Name: WDR33

Submitted by: Anonymous



Host:Rabbit

Target Name:WDR33

Sample Tissue:Human HepG2

Antibody Dilution: 1.0µg/mlWDR33 is supported by BioGPS gene expression data to be expressed in HepG2

WDR33 antibody - middle region - References

Ewing, R. M., Mol. Syst. Biol. 3, 89 (2007) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.