

Tecrl Antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12454**Specification**

Tecrl Antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8BFZ1
Other Accession	NM_153801 , NP_722496
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

Tecrl Antibody - N-terminal region - Additional Information**Gene ID** 243078**Alias Symbol** **D330017N19Rik, Srd5a2l2****Other Names**

Trans-2, 3-enoyl-CoA reductase-like, 1.3.1.-, Steroid 5-alpha-reductase 2-like 2 protein, Tecrl, Srd5a2l2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Tecrl antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Tecrl Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Tecrl Antibody - N-terminal region - Protein Information**Name** Tecrl**Synonyms** Srd5a2l2**Cellular Location**Membrane; Multi-pass membrane protein. Endoplasmic reticulum
{ECO:0000250|UniProtKB:Q5HYJ1}**Tissue Location**


Expression is highest in the heart with very low to almost undetectable levels in brain, skeletal muscle, stomach, pancreas, liver, kidney, small intestine, and uterus

Tecrl Antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Tecrl Antibody - N-terminal region - Images



Host: Rabbit
Target Name: Tecrl
Sample Tissue: Mouse Testis lysates
Antibody Dilution: 1.0µg/ml