

AGPAT2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12464**Specification**

AGPAT2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5VUD3
Other Accession	NM_001012727 , NP_001012745
Reactivity	Human, Pig
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27kDa KDa

AGPAT2 antibody - C-terminal region - Additional Information

Alias Symbol	1-AGPAT2, BSCL, BSCL1, LPAAB, LPAAT-beta
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-AGPAT2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


AGPAT2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

AGPAT2 antibody - C-terminal region - Protein Information**AGPAT2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AGPAT2 antibody - C-terminal region - Images



WB Suggested Anti-AGPAT2 Antibody Titration: 5.0µg/ml
Positive Control: Jurkat cell lysate

AGPAT2 antibody - C-terminal region - References

Niesporek, S., (2005) Br. J. Cancer 92(9), 1729-1736 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.