

MPG antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12480**Specification**

MPG antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5J9I4
Other Accession	NM_001015052 , NP_001015052
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa kDa

MPG antibody - C-terminal region - Additional Information

Alias Symbol	AAG, APNG, CRA36.1, MDG, Mid1, PIG11, PIG16, anpg, ADPG
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MPG antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


MPG antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

MPG antibody - C-terminal region - Protein Information**MPG antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MPG antibody - C-terminal region - Images



WB Suggested Anti-MPG Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate



WB Suggested Anti-MPG Antibody
Positive Control: Lane1: 50ug RCC4 lysate, Lane2: 50ug RCC4 lysate, Lane3: 50ug 786-0 lysate, Lane4: 50ug 786-0 lysate
Primary Antibody Dilution : 1:1000
Secondary Antibody : Anti-rabbit-Alexa 680
Secondary Antibody Dilution : 1:5000
Submitted by: Dr Ester Hammond, Gray Institute for Radiation Oncology and Biology-Department of Oncology, University of Oxford

MPG antibody - C-terminal region - References

Oh, J.H., (2005) Mamm. Genome 16(12), 942-954 Reconstitution and Storage: For short term use, store at 2-8 $^{\circ}$ C up to 1 week. For long term storage, store at -20 $^{\circ}$ C in small aliquots to prevent freeze-thaw cycles.