

PODXL antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12485**Specification**

PODXL antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q53ER6
Other Accession	NM_005397 , NP_005388
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53kDa KDa

PODXL antibody - N-terminal region - Additional Information

Alias Symbol **Gp200, MGC138240, PCLP, PC, PCLP-1**
Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-PODXL antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


PODXL antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PODXL antibody - N-terminal region - Protein Information**PODXL antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PODXL antibody - N-terminal region - Images



WB Suggested Anti-PODXL Antibody Titration: 5.0µg/ml
Positive Control: HepG2 cell lysate

PODXL antibody - N-terminal region - References

Casey, G., (2006) Hum. Mol. Genet. 15(5), 735-741
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.