

NRCAM antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12504**Specification**

NRCAM antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O4KMQ7
Other Accession	NM_001037132 , AAH98401
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84kDa kDa

NRCAM antibody - N-terminal region - Additional Information

Alias Symbol **KIAA0343, MGC138845, MGC138846**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NRCAM antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NRCAM antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NRCAM antibody - N-terminal region - Protein Information**NRCAM antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NRCAM antibody - N-terminal region - Images



WB Suggested Anti-NRCAM Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate