

Hepacam2 Antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12514**Specification**

Hepacam2 Antibody - N-terminal region - Product Information

Application	WB
Primary Accession	B5DEN8
Other Accession	NM_001106580 , NP_001100050
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50kDa KDa

Hepacam2 Antibody - N-terminal region - Additional Information

Alias Symbol **RGD1305697**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Hepacam2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Hepacam2 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Hepacam2 Antibody - N-terminal region - Protein Information**Hepacam2 Antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hepacam2 Antibody - N-terminal region - Images



WB Suggested Anti-Hepacam2 Antibody Titration: 1.0 µg/ml
Positive Control: Rat Small Intestine