

FCER1A antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12554

Specification

FCER1A antibody - middle region - Product Information

Application	WB
Primary Accession	P12319
Other Accession	NM_002001 , NP_001992
Reactivity	Human, Rat, Rabbit
Predicted	Human, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27kDa KDa

FCER1A antibody - middle region - Additional Information

Gene ID 2205

Alias Symbol **FCE1A, FcERI**

Other Names

High affinity immunoglobulin epsilon receptor subunit alpha, Fc-epsilon RI-alpha, FcERI, IgE Fc receptor subunit alpha, FCER1A, FCE1A

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-FCER1A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

FCER1A antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

FCER1A antibody - middle region - Protein Information

Name FCER1A

Synonyms FCE1A

Function

High-affinity receptor for immunoglobulin epsilon/IgE. Mediates IgE effector functions in myeloid cells. Upon IgE binding and antigen/allergen cross-linking initiates signaling pathways that lead to myeloid cell activation and differentiation. On mast cells, basophils and eosinophils stimulates the secretion of vasoactive amines, lipid mediators and cytokines that contribute to inflammatory response, tissue remodeling and cytotoxicity against microbes. Triggers the immediate hypersensitivity response to allergens as a host defense mechanism against helminth parasites,

pathogenic bacteria and venom toxicity. When dysregulated, it can elicit harmful life-threatening allergic and anaphylactic reactions.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Expressed in eosinophils.

FCER1A antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FCER1A antibody - middle region - Images



WB Suggested Anti-FCER1A Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Human Muscle