

NDST2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12582**Specification**

NDST2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	J3KNC8
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83kDa KDa

NDST2 Antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti-NDST2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

NDST2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NDST2 Antibody - C-terminal region - Protein Information**NDST2 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDST2 Antibody - C-terminal region - Images

168 kDa__
144 kDa__
90 kDa__
65 kDa__
40 kDa__



Host: Rabbit
Target Name: NDST2
Sample Tissue: HepG2 Cell Lysate
Antibody Dilution: 1.0µg/ml

Host: Rabbit
Target Name: NDST2
Sample Tissue: HepG2 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml