

RORC antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12593**Specification**

RORC antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q5SZR9
Other Accession	NM_001001523 , NP_001001523
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56kDa KDa

RORC antibody - N-terminal region - Additional Information

Alias Symbol	MGC129539, NR1F3, RORG, RZRG, TOR, RZR-GAMMA
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RORC antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


RORC antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RORC antibody - N-terminal region - Protein Information**RORC antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RORC antibody - N-terminal region - Images



WB Suggested Anti-RORC Antibody Titration: 1 µg/ml
Positive Control: HepG2 cell lysate

RORC antibody - N-terminal region - References

Huang, Z., (2007) Expert Opin. Ther. Targets 11(6), 737-743. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.