

Adam23 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12671**Specification**

Adam23 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9R1V7
Other Accession	NM_011780 , NP_035910
Reactivity	Human, Mouse, Rabbit, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa

Adam23 antibody - C-terminal region - Additional Information**Gene ID** 23792**Alias Symbol** **AW046396, MDC3****Other Names**

Disintegrin and metalloproteinase domain-containing protein 23, ADAM 23, Metalloproteinase-like, disintegrin-like, and cysteine-rich protein 3, MDC-3, Adam23, Mdc3

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Adam23 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Adam23 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Adam23 antibody - C-terminal region - Protein Information**Name** Adam23**Synonyms** Mdc3**Function**

May play a role in cell-cell and cell-matrix interactions. This is a non-catalytic metalloprotease-like protein (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein


Tissue Location

Brain specific..

Adam23 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Adam23 antibody - C-terminal region - Images

WB Suggested Anti-Adam23 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Mouse Heart