

Endogl1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12694**Specification**

Endogl1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8C163
Other Accession	NM_172456 , NP_766044
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

Endogl1 antibody - N-terminal region - Additional Information**Gene ID** 208194**Alias Symbol** AW557704, ENGL-B, ENGL-a, Endogl1, Endogl2, Engl, Engla, Englb**Other Names**

Nuclease EXOG, mitochondrial, 3.1.30.-, Endonuclease G-like 1, Endo G-like 1, Exog, Endogl1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Endogl1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Endogl1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Endogl1 antibody - N-terminal region - Protein Information**Name** Exog**Synonyms** Endogl1**Function**

Endo/exonuclease with nicking activity towards supercoiled DNA, a preference for single-stranded DNA and 5'-3' exonuclease activity.

Cellular Location

Mitochondrion inner membrane.

Endogl1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Endogl1 antibody - N-terminal region - Images



WB Suggested Anti-Endogl1 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Kidney