

**DDOST antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI12696****Specification**

---

**DDOST antibody - N-terminal region - Product Information**

|                   |                                                                                  |
|-------------------|----------------------------------------------------------------------------------|
| Application       | WB                                                                               |
| Primary Accession | <a href="#">Q5VWA5</a>                                                           |
| Other Accession   | <a href="#">NM_005216</a> , <a href="#">NP_005207</a>                            |
| Reactivity        | Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog |
| Predicted         | Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Horse, Bovine, Guinea Pig    |
| Host              | Rabbit                                                                           |
| Clonality         | Polyclonal                                                                       |
| Calculated MW     | 46kDa KDa                                                                        |

**DDOST antibody - N-terminal region - Additional Information**

|              |                                                                           |
|--------------|---------------------------------------------------------------------------|
| Alias Symbol | AGE-R1, KIAA0115, MGC2191, OK/SW-cl.45, OST, OST48, WBP1, AGER1, OKSWcl45 |
|--------------|---------------------------------------------------------------------------|

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 100 ul of distilled water. Final anti-DDOST antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**


DDOST antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**DDOST antibody - N-terminal region - Protein Information****DDOST antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## DDOST antibody - N-terminal region - Images



WB Suggested Anti-DDOST Antibody Titration: 5.0 µg/ml  
Positive Control: Jurkat cell lysate

## DDOST antibody - N-terminal region - References

Shibatani, T., (2005) Biochemistry 44(16), 5982-5992. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.