

**RTN2 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI12700****Specification**

---

**RTN2 antibody - N-terminal region - Product Information**

|                   |                                                                |
|-------------------|----------------------------------------------------------------|
| Application       | IHC, WB                                                        |
| Primary Accession | <a href="#">A8K7F2</a>                                         |
| Other Accession   | <a href="#">NM_206900</a> , <a href="#">NP_996783</a>          |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Predicted         | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                                         |
| Clonality         | Polyclonal                                                     |
| Calculated MW     | 51kDa KDa                                                      |

**RTN2 antibody - N-terminal region - Additional Information**

Alias Symbol **NSP2, NSPL1**

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 100 ul of distilled water. Final anti-RTN2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**


RTN2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**RTN2 antibody - N-terminal region - Protein Information****RTN2 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**RTN2 antibody - N-terminal region - Images**



Human Liver



WB Suggested Antibody Titration: 2.5 µg/ml

Positive Control: Jurkat

#### **RTN2 antibody - N-terminal region - References**

Oertle, T., (2003) FASEBJ. 17(10), 1238-1247. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.