

Bace1 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12723**Specification****Bace1 antibody - middle region - Product Information**

Application	WB
Primary Accession	P56818
Other Accession	NM_011792 , NP_035922
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Horse, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56kDa KDa

Bace1 antibody - middle region - Additional Information**Gene ID** 23821**Alias Symbol** **C76936****Other Names**

Beta-secretase 1, 3.4.23.46, Aspartyl protease 2, ASP2, Asp 2, Beta-site amyloid precursor protein cleaving enzyme 1, Beta-site APP cleaving enzyme 1, Memapsin-2, Membrane-associated aspartic protease 2, Bace1, Bace

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Bace1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Bace1 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Bace1 antibody - middle region - Protein Information**Name** Bace1 {ECO:0000312|MGI:MGI:1346542}**Synonyms** Bace**Function**

Responsible for the proteolytic processing of the amyloid precursor protein (APP) (PubMed:29325091). Cleaves at the N-terminus of the A-beta peptide sequence, between residues 671 and 672 of APP, leads to the

generation and extracellular release of beta-cleaved soluble APP, and a corresponding cell-associated C-terminal fragment which is later released by gamma-secretase (PubMed:29325091). Cleaves CHL1 (PubMed:29325091).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P56817}; Single-pass type I membrane protein. Golgi apparatus, trans-Golgi network. Endoplasmic reticulum {ECO:0000250|UniProtKB:P56817}. Endosome. Late endosome. Early endosome. Cell surface. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:P56817}. Membrane raft. Lysosome Recycling endosome. Cell projection, axon. Cell projection, dendrite. Note=Predominantly localized to the later Golgi/trans-Golgi network (TGN) and minimally detectable in the early Golgi compartments. A small portion is also found in the endoplasmic reticulum, endosomes and on the cell surface (By similarity). Colocalization with APP in early endosomes is due to addition of bisecting N-acetylglucosamine which blocks targeting to late endosomes and lysosomes (PubMed:25592972). Retrogradely transported from endosomal compartments to the trans-Golgi network in a phosphorylation- and GGA1- dependent manner (By similarity) {ECO:0000250|UniProtKB:P56817, ECO:0000269|PubMed:25592972}

Tissue Location


Expressed in the brain, specifically in neurons and astrocytes (at protein level).

Bace1 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bace1 antibody - middle region - Images



WB Suggested Anti-Bace1 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:1562500
Positive Control: Mouse Heart