

PILRB antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12729**Specification**

PILRB antibody - middle region - Product Information

Application	WB
Primary Accession	O9UKJ0
Other Accession	NM_013440 , NP_038468
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23kDa KDa

PILRB antibody - middle region - Additional Information**Gene ID** 29990**Alias Symbol** **FDFACT1, FDFACT2****Other Names**

Paired immunoglobulin-like type 2 receptor beta, Activating receptor PILR-beta, Cell surface receptor FDFACT, PILRB, FDFACT

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PILRB antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PILRB antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

PILRB antibody - middle region - Protein Information**Name** PILRB**Synonyms** FDFACT**Function**

Paired receptors consist of highly related activating and inhibitory receptors and are widely involved in the regulation of the immune system. PILRB is thought to act as a cellular signaling activating receptor that associates with ITAM-bearing adapter molecules on the cell surface.

Cellular Location


Membrane; Single-pass type I membrane protein

PILRB antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PILRB antibody - middle region - Images



WB Suggested Anti-PILRB Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: Human Muscle

PILRB antibody - middle region - References

Wilson, M.D., (2006) *Physiol. Genomics* 27(3), 201-218. Reconstitution and Storage: For short term use, store at -2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.