

KIAA0247 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12746**Specification**

KIAA0247 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O92537
Other Accession	NM_014734 , NP_055549
Reactivity	Human, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 32kDa KDa

KIAA0247 antibody - N-terminal region - Additional Information**Gene ID** 9766**Other Names**

Uncharacterized protein KIAA0247, KIAA0247

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-KIAA0247 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

KIAA0247 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

KIAA0247 antibody - N-terminal region - Protein Information**Name** SUSD6 ([HGNC:19956](#))**Function**

May play a role in growth-suppressive activity and cell death (PubMed:24652652). May be involved in the production of chemokine molecules in umbilical vein endothelial cells (HUVECs) cultured in THP1 monocyte LPS-induced medium (PubMed:20236627). Plays a role in preventing tumor onset (By similarity).

Cellular Location


Membrane; Single-pass type I membrane protein

KIAA0247 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KIAA0247 antibody - N-terminal region - Images



WB Suggested Anti-KIAA0247 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Human Liver

KIAA0247 antibody - N-terminal region - References

Gerhard, D.S., Genome Res. 14(10B), 2121-2127 (2004) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.