

RHBDL2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12781**Specification**

RHBDL2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9NX52
Other Accession	NM_017821 , NP_060291
Reactivity	Human, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Rabbit, Pig, Horse, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 34kDa kDa

RHBDL2 antibody - N-terminal region - Additional Information**Gene ID** 54933**Alias Symbol** MGC16997, RRP2**Other Names**

Rhomboid-related protein 2, RRP2, 3.4.21.105, Rhomboid-like protein 2, Rhomboid-related protein 2, N-terminal fragment, NTF, Rhomboid-related protein 2, C-terminal fragment, CTF, RHBDL2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RHBDL2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

RHBDL2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RHBDL2 antibody - N-terminal region - Protein Information**Name** RHBDL2**Function**

Involved in regulated intramembrane proteolysis and the subsequent release of functional polypeptides from their membrane anchors. Known substrate: EFN3.

Cellular Location


[Rhomboid-related protein 2, C-terminal fragment]: Cell membrane; Multi-pass membrane protein

RHBDL2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RHBDL2 antibody - N-terminal region - Images



WB Suggested Anti-RHBDL2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: PANC1 cell lysate

RHBDL2 antibody - N-terminal region - References

Gregory, S.G., (2006) Nature 441(7091), 315-321 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.