

**KIAA0515 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI12862****Specification**

---

**KIAA0515 antibody - N-terminal region - Product Information**

|                   |                                                           |
|-------------------|-----------------------------------------------------------|
| Application       | WB                                                        |
| Primary Accession | <a href="#">Q5JSZ5</a>                                    |
| Other Accession   | <a href="#">EAW87968</a>                                  |
| Reactivity        | Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog |
| Predicted Host    | Human, Mouse, Chicken, Bovine, Dog                        |
| Clonality         | Rabbit                                                    |
| Calculated MW     | Polyclonal<br>48kDa KDa                                   |

**KIAA0515 antibody - N-terminal region - Additional Information****Gene ID** 84726**Alias Symbol** RP11-334J6.1, DKFZp781F05101, DKFZp781K12107, LQFBS-1, MGC10526, BAT2L, BAT2L1, KIAA0515**Other Names**

Protein PRRC2B, HLA-B-associated transcript 2-like 1, Proline-rich coiled-coil protein 2B, PRRC2B, BAT2L, BAT2L1, KIAA0515

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-KIAA0515 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**


KIAA0515 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**KIAA0515 antibody - N-terminal region - Protein Information****Name** PRRC2B**Synonyms** BAT2L, BAT2L1, KIAA0515**KIAA0515 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**KIAA0515 antibody - N-terminal region - Images**



WB Suggested Anti-KIAA0515 Antibody Titration: 0.2-1 µg/ml

Positive Control: HepG2 cell lysate

PRRC2B is supported by BioGPS gene expression data to be expressed in HepG2