

C19orf47 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12908**Specification**

C19orf47 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8N9M1-2
Other Accession	NM_178830 , NP_849152
Reactivity	Human, Mouse, Rat, Rabbit, Guinea Pig, Dog
Predicted	Rabbit, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

C19orf47 antibody - C-terminal region - Additional Information

Alias Symbol **DKFZp686P05129, FLJ36888**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C19orf47 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


C19orf47 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

C19orf47 antibody - C-terminal region - Protein Information**C19orf47 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C19orf47 antibody - C-terminal region - Images



WB Suggested Anti-C19orf47 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate

C19orf47 antibody - C-terminal region - References

Mehrle, A., Nucleic Acids Res. 34 (DATABASE ISSUE), D415-D418 (2006) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.