

ACCSL antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12914**Specification**

ACCSL antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q4AC99
Other Accession	NM_001031854 , NP_001027025
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65kDa KDa

ACCSL antibody - N-terminal region - Additional Information**Gene ID** 390110**Other Names**

Probable inactive 1-aminocyclopropane-1-carboxylate synthase-like protein 2, ACC synthase-like protein 2, ACCSL

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ACCSL antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ACCSL antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


ACCSL antibody - N-terminal region - Protein Information**Name** ACCSL**ACCSL antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ACCSL antibody - N-terminal region - Images



WB Suggested Anti-ACCSL Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate