

Gif Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12928**Specification**

Gif Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P52787
Other Accession	NM_008118 , NP_032144
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45kDa KDa

Gif Antibody - C-terminal region - Additional Information**Gene ID** 14603**Alias Symbol** **AV073125****Other Names**

Gastric intrinsic factor, Intrinsic factor, IF, INF, Gif

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Gif antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Gif Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Gif Antibody - C-terminal region - Protein Information**Name** Cblif**Synonyms** Gif {ECO:0000312|MGI:MGI:1202394}**Function**

Promotes absorption of the essential vitamin cobalamin (Cbl) in the ileum. After interaction with CUBN, the CBLIF-cobalamin complex is internalized via receptor-mediated endocytosis (By similarity).

Cellular Location

Secreted.


Tissue Location

Gastric mucosa.

Gif Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Gif Antibody - C-terminal region - Images

Host: Rabbit

Target Name: Gif

Sample Tissue: Mouse Stomach lysates

Antibody Dilution: 1.0µg/ml