

PECI antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12932**Specification**

PECI antibody - middle region - Product Information

Application	WB
Primary Accession	Q7L124
Other Accession	NM_006117 , NP_006108
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Zebrafish, Pig, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

PECI antibody - middle region - Additional Information

Alias Symbol	ACBD2, DRS1, HCA88, dJ1013A10.3, PECI, DRS-1
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PECI antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


PECI antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

PECI antibody - middle region - Protein Information**PECI antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PECI antibody - middle region - Images



WB Suggested Anti-PECI Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

ECI2 is supported by BioGPS gene expression data to be expressed in HepG2

PECI antibody - middle region - References

Feng, X., (2004) Blood 104(8), 2425-2431 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.