

GGTL3 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12990**Specification**

GGTL3 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	B5MD39
Other Accession	NM_178026 , NP_821158
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Mouse, Rat, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 70kDa KDa

GGTL3 antibody - C-terminal region - Additional Information

Alias Symbol	D20S101, GGTL5, dJ18C9.2, GGT4, GGTL3
Other Names	
Putative gamma-glutamyltransferase light chain 3, GGTL3	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GGTL3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

GGTL3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GGTL3 antibody - C-terminal region - Protein Information

Name GGTL3


GGTL3 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GGTL3 antibody - C-terminal region - Images



WB Suggested Anti-GGTL3 Antibody Titration: 0.2-1 µg/ml
Positive Control: Human brain

GGTL3 antibody - C-terminal region - References

Dunham I., et al. Nature 402:489-495(1999).
Heisterkamp N., et al. Hum. Genet. 123:321-332(2008).