

Mettl10 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12999**Specification**

Mettl10 antibody - middle region - Product Information

Application	WB
Primary Accession	O9D853
Other Accession	NM_028095 , NP_082371
Reactivity	Human, Mouse, Rat, Rabbit, Guinea Pig, Dog
Predicted Host	Human, Rat, Rabbit, Pig, Chicken
Clonality	Rabbit
Calculated MW	Polyclonal 27kDa kDa

Mettl10 antibody - middle region - Additional Information**Gene ID** 72096**Alias Symbol** 2010208K18Rik, MGC144343**Other Names**

Methyltransferase-like protein 10, 2.1.1.-, Mettl10

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Mettl10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Mettl10 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Mettl10 antibody - middle region - Protein Information**Name** Eef1akmt2 {ECO:0000255|HAMAP-Rule:MF_03188}**Function**

Protein-lysine methyltransferase that selectively catalyzes the trimethylation of EEf1A at 'Lys-318'.

Cellular Location


Cytoplasm {ECO:0000255|HAMAP-Rule:MF_03188}. Nucleus {ECO:0000255|HAMAP-Rule:MF_03188}

Mettl10 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mettl10 antibody - middle region - Images



WB Suggested Anti-Mettl10 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Spleen

Mettl10 antibody - middle region - References

Carninci P., et al. Science 309:1559-1563(2005).