

GGT2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13000**Specification**

GGT2 antibody - N-terminal region - Product Information

Application	WB
Other Accession	XM_001129425 , XP_001129425
Reactivity	Human, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human
Clonality	Rabbit
Calculated MW	Polyclonal 62kDa KDa

GGT2 antibody - N-terminal region - Additional Information

Alias Symbol **GGT, GGT 2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GGT2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

GGT2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GGT2 antibody - N-terminal region - Protein Information**GGT2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GGT2 antibody - N-terminal region - Images



90 kDa__
65 kDa__
40 kDa__
31 kDa__
22 kDa__

WB Suggested Anti-GGT2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Hela cell lysate