

DGAT2L7 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13011**Specification**

DGAT2L7 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6IED9
Other Accession	CAD91637
Reactivity	Human, Yeast
Predicted	Human, Yeast
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24kDa KDa

DGAT2L7 antibody - C-terminal region - Additional Information

Alias Symbol **DGAT2L7,**
Other Names
Putative diacylglycerol O-acyltransferase 2-like protein DGAT2L7P, 2.3.1.-, Diacylglycerol O-acyltransferase 2-like 7 pseudogene, DGAT2L7P

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DGAT2L7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


DGAT2L7 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DGAT2L7 antibody - C-terminal region - Protein Information**DGAT2L7 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

DGAT2L7 antibody - C-terminal region - Images

WB Suggested Anti-DGAT2L7 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:1562500

Positive Control: COLO205 cell lysate