

MDM1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13033**Specification**

MDM1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	B2RB22
Other Accession	NM_020128 , NP_064513
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25kDa KDa

MDM1 antibody - N-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MDM1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


MDM1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

MDM1 antibody - N-terminal region - Protein Information**MDM1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MDM1 antibody - N-terminal region - Images



WB Suggested Anti-MDM1 Antibody Titration: 0.2-1 μ g/ml

Positive Control: Jurkat cell lysate

MDM1 is supported by BioGPS gene expression data to be expressed in Jurkat