

Mfsd11 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13087**Specification**

Mfsd11 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8BJ51
Other Accession	NM_178620 , NP_848735
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49kDa KDa

Mfsd11 antibody - C-terminal region - Additional Information**Gene ID** 69900**Alias Symbol** 2600014M03Rik, MGC113819**Other Names**

UNC93-like protein MFSD11, Major facilitator superfamily domain-containing protein 11, Protein ET, Mfsd11, Et

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Mfsd11 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Mfsd11 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Mfsd11 antibody - C-terminal region - Protein Information**Name** Mfsd11**Synonyms** Et**Cellular Location**

Membrane; Multi-pass membrane protein

Tissue Location


Widely expressed..

Mfsd11 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mfsd11 antibody - C-terminal region - Images



WB Suggested Anti-Mfsd11 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Thymus

Mfsd11 antibody - C-terminal region - References

Carninci P., et al. Science 309:1559-1563(2005).
Church D.M., et al. PLoS Biol. 7:E1000112-E1000112(2009).
Sureau A., et al. Nucleic Acids Res. 25:4513-4522(1997).
Trost M., et al. Immunity 30:143-154(2009).