

ACBD4 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13096**Specification**

ACBD4 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8NC06
Other Accession	NM_024722 , NP_078998
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35kDa KDa

ACBD4 antibody - N-terminal region - Additional Information**Gene ID** 79777

Alias Symbol	FLJ13322, FLJ90623, HMFT0700
Other Names	
Acyl-CoA-binding domain-containing protein 4, ACBD4	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ACBD4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ACBD4 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ACBD4 antibody - N-terminal region - Protein Information

Name ACBD4

Function


Binds medium- and long-chain acyl-CoA esters and may function as an intracellular carrier of acyl-CoA esters.

ACBD4 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ACBD4 antibody - N-terminal region - Images



WB Suggested Anti-ACBD4 Antibody Titration: 0.2-1 μ g/ml
Positive Control: MCF7 cell lysate

ACBD4 antibody - N-terminal region - References

Yamada S., et al. Oncogene 23:5901-5911(2004).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.