

FTHL17 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13114**Specification**

FTHL17 antibody - middle region - Product Information

Application	WB
Primary Accession	Q9BXU8
Other Accession	NM_031894 , NP_114100
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21kDa KDa

FTHL17 antibody - middle region - Additional Information**Gene ID** 53940**Alias Symbol** CT38**Other Names**

Ferritin heavy polypeptide-like 17, Cancer/testis antigen 38, CT38, FTHL17

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-FTHL17 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

FTHL17 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

FTHL17 antibody - middle region - Protein Information**Name** FTHL17**Tissue Location**

Testis specific. Also expressed in several cancers.


FTHL17 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FTHL17 antibody - middle region - Images



WB Suggested Anti-FTHL17 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:312500
Positive Control: PANC1 cell lysate

FTHL17 antibody - middle region - References

Wang P.J., et al. Nat. Genet. 27:422-426(2001).
Loriot A., et al. Int. J. Cancer 105:371-376(2003).