

ZNF766 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13164**Specification**

ZNF766 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5HY98
Other Accession	NM_001010851 , NP_001010851
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

ZNF766 antibody - C-terminal region - Additional Information**Gene ID** 90321**Other Names**

Zinc finger protein 766, ZNF766

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF766 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF766 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF766 antibody - C-terminal region - Protein Information**Name** ZNF766**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF766 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF766 antibody - C-terminal region - Images



WB Suggested Anti-ZNF766 Antibody Titration: 0.2-1 µg/ml
Positive Control: Hela cell lysate

ZNF766 antibody - C-terminal region - References

Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.